



|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495112</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 08:53 AM |
| Panel Address:    | KTL               |   |                   |                      |

## HAEMATOLOGY

### HS1.1

| TestName  | Result       | Flag       | Biological Ref. Range/Unit      |
|---|--------------|------------|---------------------------------|
| <b>CBC (Complete Blood Count), Whole Blood EDTA</b>   |              |            |                                 |
| Hemoglobin (Hb)<br><small>Photometric Cyanmethemoglobin Method</small>  | <b>7.50</b>  | <b>Low</b> | 11.5-15.2 gm/dl                 |
| Erythrocyte Count (RBCs Counts)<br><small>Optical Flow Cytometry</small>  | <b>2.94</b>  | <b>Low</b> | 3.8-4.8 10 <sup>6</sup> /ul     |
| Packed Cell Volume (PCV) Hematocrit<br><small>RBC Pulse Hight Detection</small>                                     | <b>24.10</b> | <b>Low</b> | 36-46 %                         |
| Mean Corpuscular Volume (MCV)<br><small>Automated/Calculated</small>  | <b>82.00</b> | <b>Low</b> | 83 - 101 fL                     |
| Mean Corpuscular Hemoglobin (MCH)<br><small>Automated/Calculated</small>  | 25.50        | Normal     | 24-32 pg/cell                   |
| Mean Corpuscular Hb concentration (MCHC)<br><small>Automated/Calculated</small>                                     | 31.10        | Normal     | 28 - 35 g/dL                    |
| Red Blood Cell Distribution Width Coefficient of Variation (RDW-CV)<br><small>Automated/Calculated</small>          | 16.40        | Normal     | 11.7 - 17.2 %                   |
| Red Blood Cell Distribution Width Standard Deviation (RDW-SD)<br><small>Automated/Calculated</small>                | 51.90        | Normal     | 36.4 - 58.0 10 <sup>3</sup> /uL |
| <b>Platelet Count</b><br><small>Automated Optical Flow Cytometry /Manual Calculated</small>                         | 319          | Normal     | 150-410 10 <sup>3</sup> /ul     |
| Plateletcrit (PCT)<br><small>Automated Optical Flow Cytometry /Manual Calculated</small>                            | 0.29         | Normal     | 0.15 - 0.39 %                   |
| Mean Platelet Volume (MPV)<br><small>Automated/Calculated</small>   | 9.00         | Normal     | 7.0 - 12.5 %                    |
| Platelet Distribution Width (PDW)<br><small>Automated/Calculated</small>  | 12.1         | Normal     | 9.2-17.9 %                      |
| Platelet-Large Cell Count (P-LCC)<br><small>Automated/Calculated</small>  | 68.0         | Normal     |                                 |
| Platelet Larger Cell Ratio (P-LCR)<br><small>Automated/Calculated</small>   | 21.20        | Normal     | 18.5 - 68 %                     |
| <b>Total Leukocyte Count (TLC/WBC Counts)</b><br><small>Automated Optical Flow Cytometry /Manual Calculated</small> | 5.14         | Normal     | 4.0-11 10 <sup>3</sup> /uL      |
| <b>Differential Leukocyte Count (DLC)</b><br><small>Flow Cytometry/Manual/ Microscopic</small>                      |              |            |                                 |
| Neutrophils<br><small>Impedance Flow Cytometry/ Microscopy</small>  | 56.0         | Normal     | 40 - 75 %                       |
| Lymphocytes<br><small>Impedance Flow Cytometry/ Microscopy</small>  | 40.0         | Normal     | 20 - 40 %                       |
| Monocytes<br><small>Impedance Flow Cytometry/ Microscopy</small>  | 2.0          | Normal     | 2 - 10 %                        |
| Eosinophils<br><small>Impedance Flow Cytometry/ Microscopy</small>  | 2.0          | Normal     | 1 - 6 %                         |

**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864



**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)



|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495112</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 08:53 AM |
| Panel Address:    | KTL               |   |                   |                      |

### HAEMATOLOGY

#### HS1.1

| TestName   | Result      | Flag       | Biological Ref. Range/Unit     |
|--|-------------|------------|--------------------------------|
| Basophils<br><small>Impedance Flow Cytometry/ Microscopy</small> | 0.0         | Normal     | 0 - 2 %                        |
| Absolute Neutrophil Count<br><small>Automated/Calculated</small> | 2.88        | Normal     | 2.00-8.00 10 <sup>3</sup> /uL  |
| Absolute Lymphocyte Count<br><small>Automated/Calculated</small> | 2.06        | Normal     | 1.00-3.00 10 <sup>3</sup> /uL  |
| Absolute monocyte Count<br><small>Automated/Calculated</small>   | <b>0.10</b> | <b>Low</b> | 0.20 - 1.00 10 <sup>3</sup> ul |
| Absolute Eosinophils<br><small>Automated/Calculated</small>      | 0.10        | Normal     | 0.02-0.50 10 <sup>3</sup> /ul  |
| Absolute Basophils<br><small>Automated/Calculated</small>        | <b>0.00</b> | <b>Low</b> | 0.02 - 0.10 10 <sup>3</sup> ul |

**INTERPRETATION:** **Hemoglobin:** Decreases in hemoglobin occur for the same reasons as decreased RBCs. **RBCs:** The RBC count will be low with iron deficiency, blood loss, hemolysis and bone marrow suppression. Increases may be found when one moves to a higher altitude or after prolonged physical exercise, hypoxia, Polycythemia vera. **Hematocrit:** After hemorrhage or excessive intravenous fluid infusion, the hematocrit will be low. If the patient is dehydrated, the hematocrit will be increased. **MCV:** Low values indicate the cells are microcytic and are evident with conditions such as iron deficiency, lead poisoning and Thalassemias. High values indicate macrocytic cells (large cells), and are found in conditions as megaloblastic anemia, folate or Vitamin B12 deficiency, liver disease, post-splenectomy, chemotherapy or hypothyroidism. **MCH & MCHC:** Low values are associated with iron deficiency thalassemia and malnutrition. **RDW:** Elevated levels may indicate iron deficiency or other conditions with a wide distribution of various cell sizes. **Platelets:** Thrombocytopenia (reduced platelets) can cause severe bleeding. Thrombocytopenia may occur in case of aplastic anemia, drug-induced, leukemia. Elevated platelet number can be seen in following cases essential thrombocythemia, chronic leukemia, post-splenectomy, iron deficiency anemia, malignancy, chronic infection or inflammation and other conditions which may enhance platelet function are atherosclerosis, diabetes, smoking and elevated lipid and cholesterol levels. **WBCs:** Low white blood cell count (leukopenia) may be caused by a medical condition, such as an autoimmune disorder that destroys white blood cells, bone marrow problems. Certain medications also can cause white blood cell counts to drop. If your white blood cell count is higher than normal, you may have an infection or inflammation. It could indicate that you have an immune system disorder or a bone marrow disease. A high white blood cell count can also be a reaction to medication.

**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864



**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)

|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495112</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 09:20 AM |
| Panel Address:    | KTL               |   |                   |                      |

**HAEMATOLOGY**

**HS1.1**

| TestName  | Result    | Flag        | Biological Ref. Range/Unit |
|---|-----------|-------------|----------------------------|
| <b>Erythrocyte Sedimentation Rate (ESR), Whole Blood EDTA</b> |           |             |                            |
| Erythrocyte Sedimentation Rate (ESR)<br>Westergreen Method    | <b>81</b> | <b>High</b> | <20 mm/hr                  |

**CLINICAL COMMENTS:** Erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that indirectly measures the degree of inflammation present in the body. Inflammation is part of the body's immune response. It can be acute, developing rapidly after trauma, injury or infection, for example, or can occur over an extended time (chronic) with conditions such as autoimmune diseases or cancer. Moderately elevated ESR occurs with inflammation but also with anemia, infection, pregnancy, and with aging. A very high ESR usually has an obvious cause, such as a severe infection, marked by an increase in globulins, systemic vasculitis, polymyalgia rheumatica or temporal arteritis. People with multiple myeloma or Waldenstrom's macroglobulinemia (tumors that make large amounts of immunoglobulins) typically have very high ESRs even if they don't have inflammation.

**Factors increasing ESR:**

- Advanced age
- Anemia
- Pregnancy
- High fibrinogen
- Macrocytosis
- Kidney problems
- Thyroid disease
- Some cancers, such as multiple myeloma
- Infection

**Factors decreasing ESR**

- Microcytosis
- Low fibrinogen
- Polycythemia
- Marked leukocytosis



**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864




**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)



|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495111</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 08:55 AM |
| Panel Address:    | KTL               |   |                   |                      |

### BIOCHEMISTRY & IMMUNOLOGY

#### HS1.1

| TestName  | Result        | Flag        | Biological Ref. Range/Unit   |
|---|---------------|-------------|--|
| <b>Lipid (Heart Risk) Profile, Serum</b>                          |               |             |  |
| TOTAL CHOLESTEROL<br>Chod/Pap Method                              | 145.04        | Normal      | Desirable:80-200<br>Borderline High: 200 - 239<br>High > 240 mg/dL                             |
| TRIGLYCERIDES<br>Gpo/Pap Method                                   | <b>175.38</b> | <b>High</b> | Normal 150<br>Border line High 150-199<br>High 200-499<br>Very High> 500 mg/dL                 |
| HDL CHOLESTEROL<br>Direct Enzymatic Method                        | 47.30         | Normal      | 35 - 60 mg/dL  |
| LDL CHOLESTEROL<br>Direct Enzymatic Method                        | 62.66         | Normal      | Normal: 00-100,<br>Above optimal: 101-130,<br>Borderline High: 130-160,<br>High; 160-200 mg/dL |
| VERY LOW DENSITY LIPOPROTEIN (VLDL)<br>Automated/Calculated       | <b>35.08</b>  | <b>High</b> | 0.0-30.0 mg/dL   |
| TOTAL CHOLESTEROL / HDL CHOLESTEROL Ratio<br>Automated/Calculated | 3.07          | Normal      | <5.0 mg/dL   |
| LDL / HDL CHOLESTEROL Ratio<br>Automated/Calculated               | <b>1.32</b>   | <b>Low</b>  | Less than 3.50 mg/dL   |
| TOTAL LIPIDS<br>Automated/Calculated                              | 455.38        | Normal      | 400-1000 mg/dL   |

**CLINICAL COMMENTS:** Lipid Profile is the blood test useful in screening the abnormalities associated with lipids. The results of this test can assess approximate risks for cardiovascular disease (Heart attack, Heart Failure, stroke, coronary artery disease), certain forms of pancreatitis, Hypertriglyceridemia (indicative of insulin resistance) and certain genetic disorders. Total cholesterol is an estimate of all the cholesterol in the blood. Thus, higher total cholesterol may be due to high levels of HDL or high levels of LDL. So knowing the breakdown is important. High-density lipoprotein (HDL) is good cholesterol. HDL helps carry bad cholesterol out of the bloodstream and arteries. It plays a very important role in preventing clogged arteries. So, the higher the HDL number, the better. Low-density lipoprotein (LDL) is bad cholesterol. High LDL levels increase the risk of heart disease. Your actual LDL goal depends on whether or not you have existing risk factors for heart disease, such as diabetes or high blood pressure. Very Low-density lipoprotein (VLDL) is a type of bad cholesterol that contains the highest amount of triglycerides. The higher your VLDL level, the more likely you are to have a heart attack or stroke. Triglycerides are a type of blood fat that has been linked to heart disease and diabetes. If you have high triglycerides, your total cholesterol and LDL levels may be high, as well. Lifestyle plays a large role in your triglyceride level. Smoking, excessive drinking, uncontrolled diabetes, and medications such as estrogen, steroids, and some acne treatments can contribute to high triglyceride levels. Total cholesterol to HDL ratio is useful in predicting the risk of developing atherosclerosis (plaque build-up inside the arteries). **NOTE: 10-12 Hours Fasting is Mandatory for Lipid Profile. In case of the lipemic or highly turbid due to lipoproteins mainly chylomicrons, the test cannot be performed on the specimen but the patient can request for this test again after consuming a fat free diet for at least a week.**

**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864



**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)



|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495111</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 08:56 AM |
| Panel Address:    | KTL               |   |                   |                      |

## BIOCHEMISTRY & IMMUNOLOGY

### HS1.1

| TestName  | Result        | Flag        | Biological Ref. Range/Unit |
|---|---------------|-------------|----------------------------|
| <b>Liver Function Test (LFT) - Extended, Serum</b>                          |               |             |                            |
| Bilirubin Total<br><small>Mod.Jendrassik and Grof's Method</small>          | 0.66          | Normal      | 0.1-1.2 mg/dL              |
| Bilirubin Direct<br><small>DCA METHOD</small>                               | 0.18          | Normal      | 0.00 - 0.30 mg/dL          |
| Bilirubin Indirect<br><small>Calculated</small>                             | 0.48          | Normal      | 0.1 - 1.10 mg/dL           |
| Aspartate Transaminase (AST/SGOT)<br><small>Mod. IFCC Method</small>        | 32.15         | Normal      | 0-35 U/L                   |
| Alanine Amino Transferase (ALT/SGPT)<br><small>Mod. IFCC Method</small>     | 41.40         | Normal      | 3-45 U/L                   |
| SGOT/SGPT Ratio<br><small>Automated/Calculated</small>                      | 0.78          | Normal      | 0.00 - 3.50 g/dL           |
| Alkaline Phosphatase (ALP)<br><small>Mod. IFCC Method</small>               | <b>314.35</b> | <b>High</b> | 43-115 U/L                 |
| Gamma Glutamyl Transferase (GGT)<br><small>Carboxy Substrate Method</small> | 47.50         | Normal      | 5 - 64 U/L                 |
| Protein Total<br><small>Biuret Method</small>                               | 7.02          | Normal      | 6.4-8.3 g/dL               |
| Albumin<br><small>BCG (Bromo Cresol Green ) Method</small>                  | 3.81          | Normal      | 3.5-5.6 g/dL               |
| Globulin<br><small>Automated/Calculated</small>                             | 3.21          | Normal      | 2.5 - 3.8 g/dL             |
| Albumin/Globulin Ratio (A/G)<br><small>Automated/Calculated</small>         | 1.19          | Normal      | 1.00 - 2.30 g/dL           |

**CLINICAL COMMENTS:** Liver function tests can be suggested in case of hepatitis, liver cirrhosis and monitor possible side effects of medications. A variety of diseases and infections can cause acute or chronic damage to the liver, causing inflammation (hepatitis), scarring (cirrhosis), bile duct obstructions, liver tumors, and liver dysfunction. Alcohol, drugs, some herbal supplements, and toxins can also injure the liver. A significant amount of liver damage may occur before symptoms such as jaundice, dark urine, light-colored stools, itching (pruritus), nausea, fatigue, diarrhea, and unexplained weight loss or gain appear. Early detection of liver injury is essential in order to minimize damage and preserve liver function. **Alanine aminotransferase (ALT)** A very high level of ALT is frequently seen with acute hepatitis. Moderate increases may be seen with chronic hepatitis. People with blocked bile ducts, cirrhosis, and liver cancer may have ALT concentrations that are only moderately elevated or close to normal. **Aspartate aminotransferase (AST)** A very high level of AST is frequently seen with acute hepatitis. AST may be normal to moderately increased with chronic hepatitis. In people with blocked bile ducts, cirrhosis, and liver cancer, AST concentrations may be moderately increased or close to normal. When liver damage is due to alcohol, AST often increases much more than ALT (this is a pattern seen with few other liver diseases). AST is also increased after heart attacks and with muscle injury. AST is a less sensitive and less specific marker of liver injury than ALT. AST is more elevated than ALT in alcohol-induced liver injury. AST could be elevated more than ALT like: (i) alcoholic liver disease results in mitochondrial toxicity and pyridoxal phosphate, which is a co-factor for AST; (ii) Wilson disease results in subclinical haemolysis and release of AST; (iii) the presence of liver cirrhosis; once liver cirrhosis is established, AST remains higher than ALT because of destroyed sinusoidal architecture, which results in impaired clearance of AST. **Alkaline phosphatase (ALP)** may be significantly increased with obstructed bile ducts, cirrhosis, liver cancer, and also with bone disease. Albumin is often normal in liver disease but may be low due to decreased production, especially in liver cirrhosis. **Total protein (TP)** is typically normal with liver disease. **Gamma-glutamyl transferase (GGT)** test may be used to help determine the cause of an elevated ALP. Both ALP and GGT are elevated in bile duct and liver disease, but only ALP will be elevated in bone disease. Increased GGT levels are also seen with alcohol consumption and with conditions, such as congestive heart failure.

**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864



**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
PH.D (MEDICAL BIOCHEMISTRY)

|                   |                  |   |                   |                      |
|-------------------|------------------|---|-------------------|----------------------|
| Barcode No:       | 00495111         |  | Reg No.           | 168979               |
| Patient Name:     | Mrs. ANITA       |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT        |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL     |   | Report Date:      | 29-Jun-2024 10:41 AM |
| Panel Address:    | KTL              |   |                   |                      |

### BIOCHEMISTRY & IMMUNOLOGY

#### HS1.1

| TestName  | Result | Flag   | Biological Ref. Range/Unit |
|---|--------|--------|----------------------------|
| <b>KIDNEY PROFILE (KFT), Serum</b>                |        |        |                            |
| Urea<br>GLDH KINETIC METHOD                       | 85.06  | High   | 17.0-44.0 mg/dL            |
| Creatinine ,Serum<br>Enzymatic Pap Method         | 7.12   | High   | 0.5-1.2 mg/dL              |
| Uric Acid<br>Uricase/ Pap Method                  | 6.54   | High   | 2.5-6.0 mg/dL              |
| Blood Urea Nitrogen (BUN)<br>Automated/Calculated | 39.75  | High   | 6.0 - 20.0 mg/dL           |
| BUN/Creatinine Ratio<br>Automated/Calculated      | 5.58   | Normal | 0.00 - 23.0 mg/dL          |
| Urea/Creatinine Ratio<br>Automated/Calculated     | 11.95  | Normal | 0.00 - 45.0 mg/dL          |
| Calcium Total                                     | 8.71   | Normal | 8.8 - 10.5 mg/dL           |
| <b>ELECTROLYTE PROFILE ( * )</b>                  |        |        |                            |
| Sodium  | 132.60 | Low    | 135-145 mmol/L             |
| Potassium   | 5.47   | High   | 3.5-5.2 mmol/L             |
| Chloride Serum                                    | 113.50 | High   | 97-107 mmol/L              |

#### COMMENTS: KINDLY CORRELATE WITH CLINICAL FINDINGS.


**CLINICAL COMMENTS** **UREA:** High urea levels suggest poor kidney function, congestive heart failure, shock, stress, recent heart attack or severe burns; bleeding from the gastrointestinal tract; conditions that cause obstruction of urine flow or dehydration. Low urea levels can be seen in severe liver disease or malnutrition but are not used to diagnose or monitor these conditions. Low urea levels are also seen in normal pregnancy. **CREATININE:** Increases in any renal functional impairment (intrinsic renal lesions, decreased perfusion of the kidney, or obstruction of the lower urinary tract), acromegaly and hyperthyroidism. Decreases in pregnancy, muscle wasting. **URIC ACID:** Increases in case of renal failure, disseminated neoplasms, pregnancy toxemia, psoriasis, liver disease, sarcoidosis etc. Decrease is reported in Wilson's disease, Fanconi's syndrome, xanthinuria. **SODIUM:** Increases due to water loss (severe diarrhea profuse sweating, polyuria or vomiting), hypergluco- or mineralo-corticoidism, and inadequate water intake. Decreases due to intake of free water or hypotonic solutions. Dilutional hyponatremia (liver failure, cardiac failure, nephrotic syndrome, malnutrition, renal tubular abnormalities). **POTASSIUM:** Increases due to excess destruction of cells, redistribution of K<sup>+</sup> from the intra- to the extracellular compartment (crush injuries, massive haemolysis, malignant hyperpyrexia and hyperkinetic activity). Decreased renal K<sup>+</sup> excretion (acute renal failure, some cases of chronic renal failure, Addison's disease, and other sodium-depleted states). Decreases due to excess K<sup>+</sup> loss (vomiting, diarrhea, renal tubular defects, villous adenoma of the colorectum, hypercorticoidism, etc). Redistribution hypokalemia (glucose/insulin therapy, alkalosis and periodic paralysis). **CHLORIDE:** Increases in case of dehydration, acute renal failure, renal tubular acidosis, diabetes insipidus, prolonged diarrhoea, respiratory alkalosis, salicylate toxicity, hypothalamic lesions, and adrenocortical hyperfunction. Decreases in case of excessive sweating, prolonged vomiting, adrenocortical deficiency, salt-losing nephropathy, acute intermittent porphyria, various acid base disturbances, expansion of extracellular fluid volume etc. **Urea nitrogen (BUN):** Increases in case of acute & chronic intrinsic renal disease, post renal obstruction of urine, high protein intake. Decreases with high carbohydrate/low protein diets, increased anabolic demand (late pregnancy, infancy, acromegaly), malabsorption and severe liver damage. **CALCIUM:** Increases in case of malignant neoplasms (with or without bone involvement), vit-D intoxication, primary and tertiary hyperparathyroidism, sarcoidosis, Paget's disease of bone (with immobilization), thyrotoxicosis, acromegaly, diuretic phase of renal acute tubular necrosis



Dr. DEVENDRA KUMAR  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864




Dr. MOOL KUMAR  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)



|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495111</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 10:41 AM |
| Panel Address:    | KTL               |   |                   |                      |

**BIOCHEMISTRY & IMMUNOLOGY**

**HS1.1**

| TestName              | Result      | Flag        | Biological Ref. Range/Unit |
|-----------------------|-------------|-------------|----------------------------|
| <b>eGFR, Serum</b>    |             |             |                            |
| eGFR                  | <b>7.12</b> | <b>High</b> | 0.50 - 1.30 mg/dL          |
| ESTIMATED GFR BY CKD  | <b>6.89</b> | <b>Low</b>  | >60 mL/min/1.73m2          |
| ESTIMATED GFR BY MDRD | <b>7.05</b> | <b>Low</b>  | >60 mL/min/1.73m2          |

**INTERPRETATION:**

| AGE IN YEARS | GFR IN mL/min/1.73m <sup>2</sup> |
|--------------|----------------------------------|
| 20-29        | 116                              |
| 30-39        | 107                              |
| 40-49        | 99                               |
| 50-59        | 93                               |
| 60-69        | 85                               |
| >=70         | 75                               |

**NOTE:**

1. National Kidney Disease Education program recommends the use of MDRD equation to estimate or predict GFR in adults (>=20 years) with Chronic Kidney Disease (CKD)
2. MDRD equation is most accurate for GFR <=60 mL/min/1.73m<sup>2</sup>.
3. Recalculation of estimated GFR is required for African American race.

**INTERPRETATION:**

| CKD STAGE | DESCRIPTION                           | GFR (mL/min/1.73m <sup>2</sup> ) | ASSOCIATED FINDINGS                                   |
|-----------|---------------------------------------|----------------------------------|---|
| 0         | Normal kidney function                | >90                              | No proteinuria  |
| 1         | Kidney damage with normal or high GFR | >90                              | Presence of Protein, albumin, cells or casts in urine |
| 2         | Mild decrease in GFR                  | 60-89                            | -   |
| 3         | Moderate decrease in GFR              | 30-59                            | -   |
| 4         | Severe decrease in GFR                | 15-29                            | -   |
| 5         | Kidney failure                        | <15                              | -   |

**COMMENTS:**

Modification of diet in renal disease (MDRD) equation is most thoroughly validated and superior to all the other methods for estimation of GFR. It does not require weight as a variable and yields an estimated GFR normalized to 1.73m<sup>2</sup> body surface area. Using serum creatinine alone gives a poor inference of GFR because they are inversely related and effects of age, sex and race on creatinine production complicate interpretation. For African American races a modified formula is used for calculation of GFR.

**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864



**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)

|                   |                   |   |                   |                      |
|-------------------|-------------------|---|-------------------|----------------------|
| Barcode No:       | <b>00495111</b>   |  | Reg No.           | 168979               |
| Patient Name:     | <b>Mrs. ANITA</b> |   | Reg Date          | 29-Jun-2024 08:33 AM |
| Age/Sex:          | 33Y 0M 0D/Female  |   | Sample Coll. Date | 29-Jun-2024 08:33 AM |
| Referred By:      | Dr. ANKIT         |   | Sample Rec.Date   | 29-Jun-2024 08:34 AM |
| Client Code/Name: | HR023-VISHAL      |   | Report Date:      | 29-Jun-2024 09:20 AM |
| Panel Address:    | KTL               |   |                   |                      |

**IMMUNOLOGY**

**HS1.1**

| TestName   | Result       | Flag        | Biological Ref. Range/Unit   |
|--|--------------|-------------|--|
| <b>Thyroid Function Test (TFT), Serum</b>  |              |             |  |
| Tri-Iodothyronine (TT3)<br>Electro Chemiluminescence Immunoassay (ECLIA)                                   | 1.27         | Normal      | 0.80–2.0 ng/mL   |
| Thyroxine Total (TT4)<br>Electro Chemiluminescence Immunoassay (ECLIA)                                     | 6.83         | Normal      | 4.0–12 ug/dL   |
| Thyroid Stimulating Hormone (TSH) (CMIA), ULTRA SENSITIVE<br>Electro Chemiluminescence Immunoassay (ECLIA) | <b>7.410</b> | <b>High</b> | Non-Pregnant 0.38-5.33<br>pregnant female 1st trimester 0.05-3.70<br>2nd trimester 0.31-4.35<br>3rd trimester 0.41-5.18 mU/L |

**Comments:**

T3 is physiologically more active than T4 & plays an important role in maintaining euthyroidism. T3 circulates in free form (0.3 %) and in bound form (99.7%).

T4 is predominantly bound to carrier protein - thyroid binding globulin (TBG-99.9%). T4 assay aids in diagnosis of hyperthyroidism - primary or secondary hypothyroidism & thyroid hormone resistances. T4 test must also be associated with the other test of the thyroid assessment, such as TSH & T3 as well as with the clinical examination to the patient TSH levels are subject to circadian variation, reaching peak levels between 2am to 4am and at a minimum between 6pm to 10pm. The variation is of the order of 50%; hence time of the day has influence on the measured serum TSH concentrations.

Significant numbers of patients particularly those above 55 years of age have a serum TSH level between 4.68 & 10 µIU/ml. This borderline elevation may be due to presence of SUBCLINICAL HYPOTHYROIDISM. Thyroid profile and an -thyroid (an TPO & TG) antibodies examination is suggested in all such cases.

Very low serum TSH values are observed in patients who are being treated for hypothyroidism. In such patients Serum Free T3 & Free T4 estimation may also be performed.

In Pregnancy as per American Thyroid Association Reference range for TSH is as follows:-

| Level                     | Total T3(ng/ml) | Total T4(ug/dl) | TSH(uIU/ml) | Free T3(pmol/L) | Free T4(ng/dl) |
|---------------------------|-----------------|-----------------|-------------|-----------------|----------------|
| 1 <sup>st</sup> Trimester | 1.25-2.93       | 4.60-10.50      | 0.3-4.5     | 3.2-6.8         | 0.7-2.0        |
| 2 <sup>nd</sup> Trimester | 1.54-4.00       | 6.92-12.38      | 0.5-4.6     | 3.1-5.9         | 0.5-1.60       |
| 3 <sup>rd</sup> Trimester | 1.54-4.00       | 5.98-12.98      | 0.8-5.2     | 3.1-5.9         | 0.6-1.60       |

All reports must be interpreted by treating physician only.

\*\*\*\*\* End of Report \*\*\*\*\*



**Dr. DEVENDRA KUMAR**  
MBBS, DCP, MD MA  
CONSULTANT PATHOLOGIST, DMC NO-35864




**Dr. MOOL KUMAR**  
CONSULTANT BIOCHEMIST  
Ph.D (MEDICAL BIOCHEMISTRY)



9068554251

# Dr. Sumeet Singh Dang

MBBS, MD Medicine  
DM Nephrology (CMC Vellore)  
Consultant Nephrologist & Transplant Physician  
PMC : 37861



# Dr. DANG'S KIDNEY CARE

#2 Hargobind Nagar  
Opp. Govt. Press, Sirhind Road, Patiala  
M: 99156-44663 (Only For Appointment)  
E-mail: dr\_sumeet2005@yahoo.co.in

ਕਿਡਨੀ ਰੋਗਾਂ ਦੇ ਮਾਹਿਰ

9)

BP - 152/88

Pulse - 101

SpO2 - 98

Name ..... Anita ..... 36/f ..... Date ..... 16/06/24

CKD GSD / HT  
x 4 yr

HT 3/Week

T Acetazol C / Resolone 500  
1-1-1 (w/ res)

T Dytac 20 1-1-0  
2m 4m

T Amli 2.5 0-1

Q Nisega 100 mg in 100 mg  
1/week x 8

Next Visit  
21/07/24

x RBC 4000 / 10000 r/c r/c  
wcd / 50000  
Foster

HB  
Phosphate  
K/Creat