

Sample Collected At THE CURESTA HOSPITAL



PT NAME

: MR. SUNIL KUMAR SINGH

SAMPLE REGD. AT

: 01/07/2024 17:04 =

PT. AGE/SEX

: 56Y/MALE

REPORT RELEASED ON

MOBILE NO

9431272040

PATIENT UNIQUE ID NO

: 01/07/2024 17:21 : PCLS-17031

REF. BY.

: DR . SATWIK SAURAV

REPORT STAT.

: Final

BEDWARD

:/DILALYSIS

· OPD

SEROLOGY

Unit

Normal Value

Test Done

Result

HIV 1 & 2 ANTIBODIES, SCREENING TEST Sample: Serum

NON-REACTIVE

UHID NO.: TCH3848

NON-REACTIVE

HIV 1 & 2 ANTIBODIES Immunodot Assay

HIV I & II (CARD HIV Rapid test is a qualitative test used to screen for antibodies against HIV 1 and 2 viruses. As per NACO guidelines, all positive

samples should be TEST) tested by using 3 different types of kits before report is released. This is only a screening test. All samples detected reactive by the above screening test must be confirmed by using HIV Western Blot or HIV RNA tests. A non-reactive result does not exclude the possibility of exposure to or infection with HIV or the window period. NON-REACTIVE

NON-REACTIVE

HCV RAPID CARD TEST (HEPATITIS C)

HEPATITIS B SURFACE ANTIGEN (HBsAg)

Sample: Serum

NON-REACTIVE

NON-REACTIVE

HBSAG (HEPATITIS B VIRUS) Immunochromatography

Hepatitis B surface antigen (HBsAg) is the first serologic marker appearing in the serum at 6 to 16 weeks following exposure to HBV. In acute infection, HBsAg usually disappears in 1 to 2 months after the onset of symptoms. Persistence of HBsAg for more than 6 months in duration indicates development of either a chronic carrier state or chronic HBV infection.

This is only a Screening test. All reactive samples should be confirmed by confirmatory test.

-- End Of Report--

Verified By

DR. PRERNA CHOUDHARY (MD PATHOLOGY)

For Hame Gallaction: -9973127436 In case of any discrepancy due to machine error or typing error, please get it rectified immediately.